

## **RIVER VALLEY SCHOOL DISTRICT**

660 West Daley Street	≈	Spring Green, Wisconsin 53588	≈	Phone: 608-588-2551
-----------------------	---	-------------------------------	---	---------------------

523 Exhibit

## Periods of Exclusion from School with Infectious Conditions

Personal hygiene measures such as hand washing, covering the mouth and nose when coughing or sneezing, covering weeping sores, not sharing food or drinks, and not attending school when ill or suffering from diarrhea are important means of limiting the transmission of a number of common infectious conditions.

The student/staff who have been in contact with the following conditions, to be excluded from school for the periods specified.

For the purposes of these guidelines, communicable diseases include, but are not limited to, the following:

Condition	Criteria for Exclusion from School	Onsite control and prevention measures
Mumps	Exclude for 5 days after swelling onset.	Provide immunization records for exposed individuals to public health officials Good handwashing and hygiene. Avoid kissing, sharing drinks, or utensils, use proper disinfection of surfaces and toys.
Pink Eye	None, unless fever, behavior change, or unable to avoid touching the eye. Antibiotics not required for return	Good handwashing and hygiene. Avoid kissing, sharing drinks, or utensils, use proper disinfection of surfaces and toys.
Strep Throat	Exclude for 24 hours after initiation of appropriate antibiotic and fever resolved. Fever free without fever reducing medication for 24 hours	Avoid kissing, sharing drinks, or utensils; exclude infected food handlers; minimize contact with respiratory and oral secretions. Good handwashing and hygiene.
Influenza	Exclude until fever resoled for 24 hours Fever free without fever reducing	Handwashing and good personal hygiene including covering coughs and sneezes.

	modiantian for 04	
	medication for 24	
Dortuggia (whagning	hours Exclude until after 5	Defer symptometic individuals to
Pertussis (whooping		Refer symptomatic individuals to
cough)	days of antibiotic	health care provider for evaluation.
	treatment; if no	Handwashing and good personal
	antibiotic treatment,	hygiene including covering coughs and
	exclude 21 days after	sneezes
D'a wila a a	cough onset	
Diarrhea	Exclude until	3 or more loose stools in 24 hours.
	asymptomatic for 24 hours.	Outbreaks of diarrhea in group settings
	nours.	are reportable to the public health
		department.
		Good handwashing and hygiene;
		proper disinfection of tables, toys, and
	Englands and it	food preparation areas.
Gastroenteritis (viral)	Exclude until	Good handwashing and hygiene;
	asymptomatic for 24	proper disinfection of tables, toys, and
	hours for rotavirus and 48 hours with	food preparation areas. Alcohol based
		hand products are not effective. Soap
	norovirus	and water must be used.
Hepatitis A Virus	Exclude for 14 days	Consider Hepatitis A vaccine. Exposed
	after onset of	staff should no prepare meals for
	symptoms or 10 days	others.
	after onset of	Good handwashing and hygiene;
	jaundice.	proper disinfection of tables, toys, and
Pinworm Infection	None unless has	food preparation areas.
Pinworm Infection	None, unless has diarrhea.	Frequent, good handwashing,
	diarmea.	particularly by infected child and staff
		assisting in toileting; keep fingernails
		clean and short; prevent fingers in mouth.
Clostridium Difficile	Exclude until	
		Good handwashing and hygiene;
Infection (c.diff)	asymptomatic for 48	proper disinfection of tables, toys, and
	hours.	food preparation areas. Alcohol based
		hand products are not effective. Soap
Fifth Disease	None	and water must be used
Filli Disease	INOILE	Good handwashing and hygiene;
		proper disinfection of changing tables, surfaces, and toys.
Hand-foot-and –	None, unless fever	Good handwashing and hygiene;
mouth disease	present or child	proper disinfection of changing tables,
moun uiscasc	maintain hygiene or	surfaces, and toys.
	avoid close contact	surraces, and toys.
	with others.	
	Lesions must be scab	
	over before return.	
	Fever free without	
	fever reducing	
	medication for 24	
	hours	
	nouis	

T		
Impetigo	Exclude until after	Keep fingernails clean and short.
	initiation of	Good handwashing and hygiene;
	appropriate antibiotic	proper disinfection of changing tables,
	treatment and lesion	surfaces, and toys.
	are covered or	
	crusted.	
Lice	Exclude until	Avoid sharing or storing together
	treatment or removal	personal items such s headgear, combs,
	of live lice.	and clothing.
Measles (Rubeola,	Exclude for 4 days	Assess exposure risk to who are at
measles virus)	after rash onset;	risk**; provide immunization record of
	individual from day 7	exposed individuals to public health
	through day 21	officials.
	following earliest	Contacts without documented
	exposure.	immunity (2 doses of vaccine or
	**At risk (see	laboratory proof of immunity) should
	below)**	be vaccinated.
Ringworm	Exclude until	Avoid direct contact, avoid sharing
	treatment is initiated	combs, brushes, hats, and clothing.
	or lesions are covered.	Proper disinfection of surfaces and toys
		with a fungicidal agent.
Scabies	Exclude until	Itching may continue for several weeks
	treatment is complete.	following treatment. It is not an
		indicator that treatment failed.
Varicella (chicken	Exclude until lesions	Assess exposure risk to who are at
pox)	have dried and	risk**; provide immunization record of
	crusted; exclusion of	exposed individuals to public health
	at risk** usually not	officials.
	mandated, families	
	should be notified of	
	risk.	
	Fever free without	
	fever reducing	
	medication for 24	
	hours	
Meningitis (viral)	Excluded until fever	Proper disinfection of surfaces with
	has resolved for 24	soap, water, and bleach-containing
	hours.	solution; treatment of contacts not
	Fever free without	necessary, no specific treatment.
	fever reducing	
	medication for 24	
	hours	
Coronavirus (	COVID-19 is a	Wear facial mask, frequent hand
COVID-19)	respiratory illness that	washing, avoid touching others, social
	spreads person to	distance when possible, cover cough
	person. Common	and sneezes, no sharing of drinks or
	transmission are	snacks, proper and frequent
	respiratory droplets.	disinfection of surfaces and toys.
	Incubation period is	distriction of surfaces and toys.
	meduation period is	

	2-14 days. If a person	
	is diagnosed with	
	COVID-19 than	
	excluded from school	
	for 10 days plus 24	
	hours must be fever	
	free without fever	
	reducing medication	
	and symptoms have	
	resolved. If there is a	
	direct exposure, than	
1	the student or staff is	
	excluded for 10 days.	
	The affected person	
	must provide return to	
	school letter from	
	public health	
	department or primary	
	health physician.	

\*\*At Risk= Person who are not immunized, with compromised immune system, or pregnant\*\*

REVISED:May 14, 2015APPROVED:June 15, 2015APPROVED:August 13, 2020REVISED:January 14, 2021APPROVED:February 11, 2021